

WEST[Help](#)[Logout](#)[Interrupt](#)[Main Menu](#)[Search Form](#)[Posting Counts](#)[Show S Numbers](#)[Edit S Numbers](#)[Preferences](#)[Cases](#)**Search Results -**

Term	Documents
(1 AND 2).USPT,PGPB,JPAB,EPAB,DWPI,TDBD.	0
(L1 AND L2).USPT,PGPB,JPAB,EPAB,DWPI,TDBD.	0

Database:

[US Patents Full-Text Database](#)
[US Pre-Grant Publication Full-Text Database](#)
[JPO Abstracts Database](#)
[EPO Abstracts Database](#)
[Derwent World Patents Index](#)
[IBM Technical Disclosure Bulletins](#)

Search:

L3

[Refine Search](#)[Recall Text](#)[Clear](#)**Search History**
DATE: **Tuesday, November 25, 2003** [Printable Copy](#) [Create Case](#)

<u>Set Name</u> side by side	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u> result set
<i>DB=USPT,PGPB,JPAB,EPAB,DWPI,TDBD; PLUR=YES; OP=ADJ</i>			
<u>L3</u>	11 and 12	0	<u>L3</u>
<u>L2</u>	atherosclerosis	765	<u>L2</u>
<u>L1</u>	hibiscus sabdariffa	48	<u>L1</u>

END OF SEARCH HISTORY

.L9 ANSWER 4 OF 4 CA COPYRIGHT 2003 ACS on STN
AB The effects of kenaf (*Hibiscus cannabinus*) seeds on serum and liver lipid concn. in rats were investigated. Rats were fed a hypercholesterolemic diet (contg. 1% cholesterol and 0.25% cholic acid) supplemented with 5% kenaf seed powder or 5% hot-water ext. from the calyx of Thai kenaf (*Hibiscus sabdariffa*). The serum cholesterol concn. was significantly lower in the rats fed the diet supplemented with seed powder than in rats of the control group. The ext. from the flower did not exert a similar effect. The HDL-cholesterol concn. in the serum was not converted, but the (VLDL+LDL)-cholesterol concn. decreased as compared to the control group. No significant differences in liver cholesterol concn. were obsd. for all substances tested. Rats were fed a diet contg. polysaccharide extd. from the seeds collected in the Hunan and Zhejiang regions of China. The serum cholesterol concn. was significantly lower in the rats fed a diet supplemented with 1% or 5% polysaccharides as compared to control group, but did not differ significantly between the two regional seed sources. There was no significant difference in liver cholesterol concn. between the control and the treatment groups. The results suggest that the polysaccharides extd. from kenaf seeds have serum cholesterol-lowering effects in rats.
AN 135:221097 CA
TI Cholesterol-lowering effect of water-soluble polysaccharides from kenaf (*Hibiscus cannabinus*) seeds in rats I
AU Tamaki, Yasutomo; Kinjo, Kazuhiko; Uechi, Shuntoku; Hongo, Fujiya; Sameshima, Kazuhiko; Yaga, Shiryo
CS Fac. Agric., Univ. Ryukyus, Okinawa, 903 0213, Japan
SO Mokuzaigakkaishi (2001), 47(2), 159-163
CODEN: MKZGA7; ISSN: 0021-4795
PB Nippon Mokuzaigakkai
DT Journal
LA Japanese